

AD611427

AD611427

MAXIMIZING A SECOND-DEGREE POLYNOMIAL
ON THE UNIT SPHERE

BY

GEORGE E. FORSYTHE and GENE H. GOLUB


COPY	2	OF	3	YMD
HARD COPY			\$.	2.00
MICROFICHE			\$.	0.50

35-p

TECHNICAL REPORT CS16

FEBRUARY 5, 1965

COMPUTER SCIENCE DEPARTMENT
School of Humanities and Sciences
STANFORD UNIVERSITY



ARCHIVE COPY

MAXIMIZING A SECOND-DEGREE POLYNOMIAL ON THE
UNIT SPHERE ^{*/}

by

George E. Forsythe

and

Gene H. Golub

ABSTRACT

Let A be a hermitian matrix of order n , and b a known vector in C^n . The problem is to determine which vectors make $\phi(x) = (x-b)^H A(x-b)$ a maximum or minimum on the unit sphere $U = \{x : x^H x = 1\}$. The problem is reduced to the determination of a finite point set, the spectrum of (A,b) . The theory reduces to the usual theory of hermitian forms when $b = 0$.

^{*/}

Reproduction in Whole or in Part is Permitted for any Purpose of the United States Government. This report was supported in part by Office of Naval Research Contract Nonr-225(37) (NR-044-211) at Stanford University.

1. The problem.

Let A be a hermitian square matrix of complex elements and order n . Let b be a known n -vector of complex numbers. For each complex n -vector x , the nonhomogeneous quadratic expression

$$(1.1) \quad \Phi(x) = (x-b)^H A (x-b)$$

(H denotes complex conjugate transpose) is a real number. The problem, suggested to us by C. R. Rao of the Indian Statistical Institute, Calcutta, is to maximize (or minimize) $\Phi(x)$ for complex x on the unit sphere $S = \{x: x^H x = 1\}$. Since Φ is a continuous function on the compact set S , such maxima and minima always exist.

In summary, our problem is:

$$(1.2) \quad \text{maximize or minimize } \Phi(x) \text{ subject to } x^H x = 1.$$

The purpose of this note is to reduce the problem (1.2) to the determination of a certain finite real point set which we shall call the spectrum of the system (A, b) (defined at end of Sec. 1), and show that a unique number λ in the spectrum determines the one or more $x = x^\lambda$ which maximize $\Phi(x)$ for given b . Theorem (4.1) is the main result. The development is an extension to general b of the familiar theory for the homogeneous case when $b = \theta$, the zero vector. No consideration to a practical computer algorithm is given here.

In Sec. 7 we show that determining the least-squares solution of an arbitrary system of linear equations $Cy = f$, subject to the quadratic

constraint $y^H y = 1$, is a special case of problem (1.2).

Phillips (9.2) and Twomey (9.3) begin the actual numerical solution of certain integral equations by approximating them with algebraic problems very closely related to the minimum problem (1.2).

Let $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ be the (necessarily real) eigenvalues of A , and let $\{u_1, \dots, u_n\}$ be a corresponding real orthonormal set of eigenvectors, with $Au_i = \lambda_i u_i$ ($i=1, \dots, n$).

Let a given b be written

$$(1.3) \quad b = \sum_{i=1}^n b_i u_i.$$

(1.4) Theorem. If x is any vector in S for which $\Phi(x)$ is stationary with respect to S , then there exists a real number $\lambda = \lambda(x)$ such that

$$(1.5) \quad A(x-b) = \lambda x,$$

$$(1.6) \quad x^H x = 1.$$

Conversely, if any real λ and vector x satisfy (1.5, 1.6), then x renders $\Phi(x)$ stationary.

Proof. Let x_0 be a point of S . Now, as shown in lemma (8.7), $\Phi(x)$ is stationary at x_0 with respect to x in S , if and only if there exists a real Lagrange multiplier λ such that $\psi(x) = (x-b)^H A(x-b) - \lambda x^H x$ is stationary at x_0 with respect to all neighboring complex vectors x .

Since

$$0 = \frac{1}{2} \text{grad } \psi(x_0) = A(x_0 - b) - \lambda x_0,$$

the theorem is proved.

To see what conditions are satisfied by the λ of theorem (1.4), we note that the system (1.5, 1.6) is equivalent to the system

$$(1.7) \quad (A - \lambda I)x = Ab, \quad ,$$

$$(1.8) \quad x^H x = 1 \quad .$$

Let

$$x = \sum_{i=1}^n x_i u_i \quad .$$

Then (1.7) is equivalent to

$$(1.9) \quad \sum_{i=1}^n (\lambda_i - \lambda) x_i u_i = \sum_{i=1}^n \lambda_i b_i u_i \quad .$$

Definition. By the spectrum of the pair (A, b) we mean the set of all real λ for which there exists an x such that (1.7) and (1.8) are satisfied.

Given any λ, x satisfying (1.7) and (1.8), we shall say that x belongs to λ , and frequently write x in the form x^λ .

Note that the spectrum of (A, θ) is the ordinary spectrum $\{\lambda_i\}$ of A .

2. Special case: no $\lambda_i b_i = 0$.

Assume for the present section that $\lambda_i b_i \neq 0$ (all i). This implies that all $\lambda_i \neq 0$, i.e., that A is nonsingular. If λ is in the spectrum of (A, b) , (1.9) implies that $\lambda \neq \lambda_i$ for all i , and also that

$$(2.1) \quad x_i = \frac{\lambda_i b_i}{\lambda_i - \lambda} \quad (i=1, \dots, n) \quad .$$

Then the requirement that

$$(2.2) \quad x^H x = \sum_{i=1}^n |x_i|^2 = 1$$

is equivalent to the condition

$$(2.3) \quad g(\lambda) = \sum_{i=1}^n \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda|^2} = 1.$$

Although all λ corresponding to stationary values of $\Phi(x)$ are known by theorem (1.4) to be real, it is useful to define $g(\lambda)$ by (2.3) for all complex λ not in $\{\lambda_i\}$.

Let G be the set of complex numbers λ such that $g(\lambda) = 1$. For small enough $\sum_{i=1}^n |b_i|$, G is the union of n simple closed curves in the complex plane, the k -th of which surrounds λ_k . As the $|b_i|$ grow, adjacent curves first coalesce in double points, and then merge into single curves. For very large values of all $|b_i|$, G is one simple closed curve including all $\{\lambda_i\}$ in its interior. The family of sets G resembles the family of lemniscates $\prod_{i=1}^n |\lambda - \lambda_i| = \text{const.}$

Note, moreover, that $g(\lambda) > 1$ for λ inside any component curve G_j of G , while $g(\lambda) < 1$ in the exterior of all components G_j of G .

Now we shall show for the special case of Sec. 2 that each λ in G determines a unique x^λ which satisfies (1.7, 1.8). For that x^λ

$$(2.4) \quad \phi(x^\lambda) = f(\lambda) \quad ,$$

where we define f by

$$(2.5) \quad f(\lambda) = |\lambda|^2 \sum_{i=1}^n \frac{\lambda_i |b_i|^2}{|\lambda_i - \lambda|^2} \quad .$$

Fix λ , and drop the superscript λ on x . To prove (2.4), note that (1.7) says $(\lambda_i - \lambda)x_i = \lambda_i b_i$. Thus

$$\begin{aligned} (\lambda_i - \lambda)(x_i - b_i) &= \lambda_i b_i - b_i(\lambda_i - \lambda) \\ &= \lambda b_i \quad . \end{aligned}$$

Hence

$$x_i - b_i = \frac{\lambda b_i}{\lambda_i - \lambda} \quad ,$$

and

$$\begin{aligned} \phi(x) &= \sum_{i=1}^n \lambda_i |x_i - b_i|^2 \\ &= |\lambda|^2 \sum_{i=1}^n \frac{\lambda_i |b_i|^2}{|\lambda_i - \lambda|^2} \\ &= f(\lambda) \quad , \end{aligned}$$

proving (2.4).

Since the Lagrange multipliers λ must be real, the spectrum of (A, b) is the intersection of G with the real axis. This consists of from

2 to $2n$ distinct real numbers. How many numbers are actually in the spectrum depends on b ; this will be discussed in Sec. 5 for $n = 2$.

We wish to determine which λ in the spectrum corresponds to the maximum [minimum] value of $f(\lambda)$. Let G_j be any component curve of the set G .

(2.6) Theorem. The maximum and minimum real parts of λ , for λ in any one G_j , both occur for λ on the real axis.

Proof. Let $\lambda = \sigma + i\tau$, with σ, τ real. Then

$$g(\lambda) = g_1(\sigma, \tau) = \sum_{i=1}^n \frac{\lambda_i^2 |b_i|^2}{(\sigma - \lambda_i)^2 + \tau^2}.$$

Hence, for $\tau > 0$ and fixed σ , $g(\lambda)$ strictly decreases as τ increases. Then in the upper half plane $\tau > 0$, any line $\sigma = \text{constant}$ intersects G_j in exactly one point. The theorem follows from this.

Definition. Let $\Lambda_R[\Lambda_L]$ denote the unique real value of λ of maximum [minimum] real part in the set G .

(2.7) Theorem. Under the assumptions that A is regular (i.e., $\lambda_i \neq 0$ for all i) and $b_i \neq 0$ ($i=1,2,\dots,n$), for all λ in G such that $\lambda \neq \Lambda_R$, $\lambda \neq \Lambda_L$ we have

$$f(\Lambda_L) < f(\lambda) < f(\Lambda_R).$$

Proof. Let $a_i = \lambda_i^2 |b_i|^2$ ($i=1,\dots,n$). Introduce two independent complex variables λ, μ , where μ will later be set equal to $\bar{\lambda}$. In order to study the gradients of the functions g, f , and h (defined below) for

complex λ , we shall use the tools of Sec. 8. This requires extending these functions into the space of λ and μ .

Let $\lambda = \sigma + i\tau$ (σ, τ real). For all complex $\lambda \neq \lambda_1$, define the functions g_1 and g_2 by

$$g(\lambda) = g_1(\sigma, \tau) = g_2(\lambda, \bar{\lambda}) ,$$

where

$$g_2(\lambda, \mu) = \sum_{i=1}^n \frac{a_i}{(\lambda_1 - \lambda)(\lambda_1 - \mu)} .$$

(This definition is consistent with (2.3).) Then

$$\frac{1}{2} \left(\frac{\partial g_1}{\partial \sigma} + i \frac{\partial g_1}{\partial \tau} \right) = \left[\frac{\partial g_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} , \quad \text{by lemma (8.1)}$$

$$\begin{aligned} (2.8) \quad &= \sum_{i=1}^n \frac{a_i}{(\lambda_1 - \lambda)(\lambda_1 - \bar{\lambda})^2} \\ &= \sum_{i=1}^n \frac{a_i}{|\lambda_1 - \lambda|^2 \cdot (\lambda_1 - \bar{\lambda})} . \end{aligned}$$

For all complex $\lambda \neq \lambda_1$, define $f(\lambda)$ by (2.5). We then define the functions f_1 and f_2 by

$$f(\lambda) = f_1(\sigma, \tau) = f_2(\lambda, \bar{\lambda}) ,$$

where

$$f_2(\lambda, \mu) = \lambda_{\mu} \sum_{i=1}^n \frac{a_i}{\lambda_i(\lambda_i - \lambda)(\lambda_i - \mu)} .$$

Then

$$\frac{1}{2} \left(\frac{\partial f_1}{\partial \sigma} + i \frac{\partial f_1}{\partial \tau} \right) = \left[\frac{\partial f_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} , \quad \text{by lemma (8.1)}$$

$$= \lambda \sum_{i=1}^n \frac{a_i}{\lambda_i |\lambda_i - \lambda|^2} + \lambda \bar{\lambda} \sum_{i=1}^n \frac{a_i}{\lambda_i (\lambda_i - \lambda)(\lambda_i - \bar{\lambda})^2}$$

$$= \lambda \sum_{i=1}^n \frac{a_i}{\lambda_i |\lambda_i - \lambda|^2} \left[1 + \frac{\bar{\lambda}}{\lambda_i - \bar{\lambda}} \right]$$

$$= \lambda \sum_{i=1}^n \frac{a_i}{|\lambda_i - \lambda|^2 (\lambda_i - \bar{\lambda})}$$

$$= \lambda \left[\frac{\partial g_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} , \quad \text{by (2.8)} .$$

i.e.,

$$(2.9) \quad \left[\frac{\partial f_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} = \lambda \cdot \left[\frac{\partial g_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} .$$

While it is possible to use (2.9) to study the behavior of $f(\lambda)$ on the set G where $g(\lambda) - 1 = 0$, it is more convenient here and in Sec. 3 to introduce a new function $h(\lambda)$, which agrees with $f(\lambda)$ on G . For all complex $\lambda \neq \lambda_i$, define

$$(2.10) \quad h(\lambda) = f(\lambda) + \frac{\lambda + \bar{\lambda}}{2} [1 - g(\lambda)] ,$$

and note that

$$(2.11) \quad h(\lambda) = f(\lambda), \quad \text{for } \lambda \in G .$$

As with f and g , we introduce functions h_1 and h_2 so that

$$h(\lambda) = h_1(\sigma, \tau) = h_2(\lambda, \mu) ,$$

where

$$h_2(\lambda, \mu) = f_2(\lambda, \mu) + \frac{\lambda + \mu}{2} [1 - g_2(\lambda, \mu)] .$$

Then

$$(2.12) \quad \begin{aligned} \frac{\partial h_2}{\partial \mu} &= \frac{\partial f_2}{\partial \mu} + \frac{1}{2} [1 - g_2(\lambda, \mu)] - \frac{\lambda + \mu}{2} \frac{\partial g_2}{\partial \mu} \\ &= \frac{\lambda - \mu}{2} \frac{\partial g_2}{\partial \mu} + \frac{1}{2} [1 - g_2(\lambda, \mu)] , \quad \text{by (2.9)} . \end{aligned}$$

Hence

$$(2.13) \quad \begin{aligned} \frac{1}{2} \left(\frac{\partial h_1}{\partial \sigma} + 1 \frac{\partial h_1}{\partial \tau} \right) &= \left[\frac{\partial h_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} , \quad \text{by (8.1)} \\ &= \frac{\lambda - \bar{\lambda}}{2} \left[\frac{\partial g_2}{\partial \mu} \right]_{\mu=\bar{\lambda}} + \frac{1}{2} [1 - g_2(\lambda, \bar{\lambda})] \\ &= \frac{\tau_1}{2} \left(\frac{\partial g_1}{\partial \sigma} + 1 \frac{\partial g_1}{\partial \tau} \right) + \frac{1}{2} [1 - g(\lambda)] . \end{aligned}$$

Now any component G_j of the set G where $g(\lambda) = 1$ encloses a region where $g(\lambda) > 1$. On G the gradient vector of g ,

$$\frac{\partial g_1}{\partial \sigma} + i \frac{\partial g_1}{\partial \tau} ,$$

is non-zero, is normal to G_j , and points to the interior of G_j . Then, by (2.12), the gradient vector of h on G_j , namely

$$\frac{\partial h_1}{\partial \sigma} + i \frac{\partial h_1}{\partial \tau} = i\tau \left(\frac{\partial g_1}{\partial \sigma} + i \frac{\partial g_1}{\partial \tau} \right) ,$$

is non-zero for $\tau \neq 0$ and points along the tangent to G_j in the direction of increasing σ . Hence

$$(2.14) \quad \begin{cases} h(\lambda) \text{ is strictly increasing, as } \lambda \text{ traces } G_j \text{ in} \\ \text{the direction of increasing } \sigma. \end{cases}$$

From (2.14) it follows that $h(\lambda)$ assumes its maximum value, for each separate component curve G_j of G , at the point β_j on G_j of maximum real part. By theorem (2.6), β_j is on the axis of real λ .

Note that setting $\mu = \bar{\lambda} = \lambda$ in (2.12) yields the result that

$$(2.15) \quad h'(\lambda) = 1 - g(\lambda), \quad \text{for real } \lambda .$$

To complete the proof of the present theorem, we must show that $f(\lambda)$ is larger at the point α_j of least real part on the component G_j of G

than it is at the right-most point β_{j-1} of the component G_{j-1} of G immediately to the left of G_j .

Note that g is continuous for $\lambda \in [\beta_{j-1}, \alpha_j]$, and that $g(\beta_{j-1}) = g(\alpha_j) = 1$ but $g(\lambda) < 1$ for $\beta_{j-1} < \lambda < \alpha_j$. Then

$$\begin{aligned} h(\alpha_j) &= h(\beta_{j-1}) + \int_{\beta_{j-1}}^{\alpha_j} h'(\lambda) d\lambda \\ &= h(\beta_{j-1}) + \int_{\beta_{j-1}}^{\alpha_j} [1-g(\lambda)] d\lambda, \quad \text{by (2.15)} \\ &> h(\beta_{j-1}), \quad \text{since } g(\lambda) < 1. \end{aligned}$$

Thus

$$(2.16) \quad h(\beta_{j-1}) < h(\alpha_j),$$

as was to be proved.

We conclude that $h(\lambda)$ increases, as λ increases along the real axis between adjacent components of G . Since $h(\lambda) = f(\lambda)$ on G , we see from (2.14) and (2.16) that

$$\max_{\lambda \in G} f(\lambda) = f(\Lambda_R),$$

$$\min_{\lambda \in G} f(\lambda) = f(\Lambda_L).$$

It follows trivially from theorem (2.7) that the maximum and minimum values of $f(\lambda)$ over the real numbers in G (i.e., over the spectrum of (A,b) in the present case) are also $f(\Lambda_R)$ and $f(\Lambda_L)$, respectively.

By (1.9), our condition that $\text{no } \lambda_i b_i = 0$ implies that $\lambda \neq \lambda_i$ for all i and for all λ in G . Hence Λ_L and Λ_R are not eigenvalues of A , and so neither $A - \Lambda_L I$ nor $A - \Lambda_R I$ is a singular matrix. Therefore we can solve equation (1.7) uniquely for x_{\max} and x_{\min} :

$$x_{\max} = x^{\Lambda_R} = (A - \Lambda_R I)^{-1} A b ,$$

$$x_{\min} = x^{\Lambda_L} = (A - \Lambda_L I)^{-1} A b .$$

These equations give unique solutions to the problem of minimizing and maximizing $\Phi(x) = (x-b)^H A (x-b)$, for nonsingular A and b such that $\text{no } b_i = 0$.

It would be desirable to be able to prove that $h(\alpha_j) < h(\beta_j)$, in the notation of theorem (2.7), without analyzing $h(\lambda)$ and $g(\lambda)$ for complex values of λ .

3. General case: Some $\lambda_i b_i = 0$.

We now study the general case where one or more $\lambda_i b_i = 0$. To be explicit, let $\mathcal{C} = \{\alpha : \lambda_\alpha b_\alpha = 0\}$, a set of integers. We wish to examine the spectrum of (A,b) .

Define \mathcal{L} as the set $\{\lambda_\alpha : \alpha \in \mathcal{C}\}$.

First, given one $\alpha \in \mathcal{C}$, if $\lambda \neq \lambda_\alpha$ and λ is in the spectrum of (A, b) with corresponding vector x^λ , then (1.9) shows that $x_\alpha^\lambda = 0$. Thus, if λ is in the spectrum but not in the set \mathcal{L} , then $x_\alpha^\lambda = 0$ for all $\alpha \in \mathcal{C}$ and, just as in the derivation of (2.3), λ will satisfy the equation

$$(3.1) \quad \sum_{i=1}^n \frac{\lambda_1^2 |b_i|^2}{|\lambda_1 - \lambda|^2} = 1.$$

Conversely, any real solution λ of (3.1) which is not in the set \mathcal{L} will be in the spectrum of (A, b) , and its corresponding vector x^λ will have $x_\alpha^\lambda = 0$ for all $\alpha \in \mathcal{C}$. If we interpret $0/0$ in (3.1) as 0 , then it is possible that some eigenvalue λ_k in \mathcal{L} will also satisfy (3.1). If so, we will show that this λ_k is also in the spectrum of (A, b) . However, the spectrum may also contain eigenvalues λ_k in \mathcal{L} which do not satisfy (3.1), as we shall now show. No eigenvalue λ_k not in \mathcal{L} can be in the spectrum, because $\lambda = \lambda_k$ would make the left side of (3.1) equal to ∞ .

Fix attention on one λ_k for $k \in \mathcal{C}$. We wish to examine the possibility that this λ_k is in the spectrum of (A, b) . Let m be the multiplicity of λ_k as an eigenvalue of A . Let $\mathcal{D}_k = \{i: \lambda_i = \lambda_k\}$ so that $\text{card}(\mathcal{D}_k) = m$. If λ_k is in the spectrum, then (1.9) shows that $\lambda_\alpha b_\alpha = 0$ for all α in \mathcal{D}_k . Moreover, if λ_k is in the spectrum, then the corresponding vector $x = x^{\lambda_k}$ has the properties

$$x_i = \frac{\lambda_1 b_i}{\lambda_1 - \lambda_k} \quad (i \notin \mathcal{D}_k),$$

and, by (2.1), also

$$(3.2) \quad \sum_{i \notin \mathcal{A}_k} |x_i|^2 = \sum_{i \notin \mathcal{A}_k} \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda_k|^2} \leq 1.$$

Conversely, if (3.2) holds then we can always define x_i for all $i \in \mathcal{A}_k$ in such a way that

$$(3.3) \quad \sum_{i \in \mathcal{A}_k} |x_i|^2 = 1 - \sum_{i \notin \mathcal{A}_k} \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda_k|^2}.$$

Hence, by (3.2) and (3.3), equation (1.8) holds and, since (1.7) is satisfied, λ_k is in the spectrum of (A, b) .

If equality holds in (3.2) then x_i must be 0 for all $i \in \mathcal{A}_k$; i.e., λ_k satisfies (3.1), and x^{λ_k} is unique. But if inequality < holds in (3.2), then there is an $(m-1)$ -dimensional sphere \mathcal{U} of values of $\{x_i\}$, for $i \in \mathcal{A}_k$, which satisfy (3.3). For, if a point $(x_{i_1}, \dots, x_{i_m})$ is in \mathcal{U} , then so are all points of form

$$(x_{i_1} e^{i\theta_1}, \dots, x_{i_m} e^{i\theta_m}) \quad (\text{all } \theta_i \text{ real}),$$

since $\sum_{i \in \mathcal{A}_k} |x_i|^2$ is constant for all of these. In this case uniqueness of x^{λ_k} is lost. The sphere is analogous to (in fact is a generalization of) the sphere of unit eigenvectors of a hermitian matrix A belonging to an eigenvalue of multiplicity m .

Note that an inequality $<$ in (3.2) states that λ_k is in the exterior of the graph

$$G = \left\{ \lambda : \sum_{i \notin \mathcal{O}_k} \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda|^2} = 1 \right\},$$

i.e., λ_k can be joined to ∞ by an arc not cutting G . Thus, in brief, the spectrum of (A, b) consists of the union of all real numbers in the set

$$(3.4) \quad G = \left\{ \lambda : \sum_{i=1}^n \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda|^2} = 1 \right\},$$

where we interpret $0/0$ as 0 , with those numbers λ_k which are exterior to the graph G . (If G is the null set, then $b = \theta$ and the spectrum of (A, θ) consists of all eigenvalues λ_k .)

We must now examine $\Phi(x^\lambda)$ for λ in the spectrum of (A, b) . The study of $\Phi(x^\lambda)$, for real $\lambda \in G$ in (3.4) is the same as in Sec. 2, and yields the same results (2.4) and (2.5): First, for $\lambda \in G$,

$$\Phi(x^\lambda) = f(\lambda) = |\lambda|^2 \sum_{i=1}^n \frac{\lambda_i |b_i|^2}{|\lambda_i - \lambda|^2}, \text{ where } 0/0 = 0. \text{ Second, let } \mu_R,$$

μ_L be the right-most [resp. left-most] points of G . Then $f(\mu_R)$ maximizes [resp. $f(\mu_L)$ minimizes] $f(\lambda)$ for $\lambda \in G$. It remains to consider $\Phi(x^{\lambda_k})$, for eigenvalues λ_k outside G .

(3.5) Theorem. For any λ in the spectrum of (A, b) we have

$$(3.6) \quad \Phi(x^\lambda) = h(\lambda) = f(\lambda) + \lambda[1-g(\lambda)] ,$$

where $f(\lambda)$ is given by (2.5), with $0/0$ interpreted as 0 .

Proof. Take any λ in the spectrum of (A, b) .

If $\lambda \neq \lambda_k$ ($k=1, \dots, n$), then $\lambda \in G$, and everything proceeds as in the proof of (2.4), showing that $\Phi(x^\lambda) = f(\lambda)$. Since $g(\lambda) = 1$, we have proved (3.6) when $\lambda \neq \lambda_k$.

If $\lambda = \lambda_k$, an eigenvalue of A , let x_i denote the i -th coordinate of any x^{λ_k} which satisfies (1.5) and (1.6) (and hence (3.2) and (3.3)).

Since λ_k is in the spectrum of (A, b) , we have $\lambda_i b_i = 0$ for all $i \in \mathcal{D}_k$, where \mathcal{D}_k is defined above after (3.1), and as $\lambda_i |x_i - b_i|^2 = \lambda_i |x_i|^2 = \lambda_k |x_i|^2$, for all $i \in \mathcal{D}_k$. Then, by (3.3),

$$(3.7) \quad \sum_{i \in \mathcal{D}_k} \lambda_i |x_i - b_i|^2 = \lambda_k \left[1 - \sum_{i \notin \mathcal{D}_k} \frac{\lambda_i |b_i|^2}{(\lambda_i - \lambda_k)^2} \right]$$

$$= \lambda_k \left[1 - \sum_{i=1}^n \frac{\lambda_i |b_i|^2}{(\lambda_i - \lambda_k)^2} \right] , \text{ where } 0/0 = 0 .$$

Moreover, like (2.4) we can prove

$$\begin{aligned}
 (3.8) \quad \sum_{1 \notin \theta_k} \lambda_1 |x_1 - b_1|^2 &= \lambda_k^2 \sum_{1 \notin \theta_k} \frac{\lambda_1 |b_1|^2}{(\lambda_1 - \lambda_k)^2} \\
 &= \lambda_k^2 \sum_{i=1}^n \frac{\lambda_1 |b_1|^2}{(\lambda_1 - \lambda_k)^2}, \quad \text{where } 0/0 = 0.
 \end{aligned}$$

Adding (3.7) to (3.8), we get

$$\begin{aligned}
 (3.9) \quad \phi(x^\lambda) &= \sum_{i=1}^n \lambda_1 |x_1 - b_1|^2 \\
 &= \lambda_k^2 \sum_{i=1}^n \frac{\lambda_1 |b_1|^2}{(\lambda_1 - \lambda_k)^2} + \lambda_k \left[1 - \sum_{i=1}^n \frac{\lambda_1 |b_1|^2}{(\lambda_1 - \lambda_k)^2} \right] \\
 &= f(\lambda_k) + \lambda_k [1 - g(\lambda_k)] \\
 &= h(\lambda_k).
 \end{aligned}$$

This proves (3.6) when $\lambda = \lambda_k$.

It is property (3.6) of h which motivated our use of h in Sec. 2.

Note. It is easily shown from (3.6) or (3.9) that, for all λ in the spectrum of (A, b) ,

$$(3.10) \quad h(\lambda) = \lambda + \lambda \sum_{i=1}^n \frac{\lambda_1 |b_1|^2}{\lambda - \lambda_1}, \quad \text{where } 0/0 = 0.$$

If λ is in the spectrum of (A, b) , but is not an eigenvalue of A , we can derive (3.10) as follows. Let x belong to λ . Then

$$\Phi(x) = (x-b)^H A(x-b)$$

$$= (x-b)^H \lambda x, \quad \text{by (1.5)}$$

$$= \lambda x^H x - \lambda b^H x$$

$$= \lambda - \lambda b^H x, \quad \text{by (1.6)}$$

$$= \lambda - \lambda b^H (A - \lambda I)^{-1} A b, \quad \text{by (1.7)}$$

$$\lambda - \lambda \sum_{i=1}^n \frac{\lambda_i |b_i|^2}{\lambda_i - \lambda}.$$

We shall not make use of (3.10) here.

We now use formula (3.6) to extend the domain of h to all real λ where $g(\lambda) < \infty$, i.e., to all λ except where, for some i , $\lambda = \lambda_i$ and $\lambda_i b_i \neq 0$.

As stated before (3.5), we know that the largest value of $\Phi(x^\lambda) = h(\lambda)$ for λ in G occurs at the right-most point μ_R of G . It remains to see whether $h(\lambda_k)$ may be still larger for any λ_k in the spectrum of (A, b) , if $\mu_R < \lambda_k$.

The answer is furnished by formula (2.15), which is valid for the general case of Sec. 3 with the understanding that $0/0 = 0$. Thus h is increasing on all segments of the real axis between or exterior to components of the curve G . It follows that $h(\lambda)$ takes its maximum at the rightmost point λ_R of the spectrum of (A, b) and its minimum value at the leftmost point λ_L of the spectrum of (A, b) , whether or not these are eigenvalues of A .

From the considerations following (3.3), we see that the maximizing x is unique if $\lambda_R \in G$. If, however, λ_R is not in G and is an eigenvalue of A of multiplicity m , then the maximizing x include all points of an $(m-1)$ -sphere of nonzero radius, whose center is not at θ when $b \neq \theta$.

The above result about λ_R and λ_L for the case where some $\lambda_i b_i = 0$ can be obtained by continuity from the case where no $\lambda_i b_i = 0$. It is not clear that we could use continuity to deduce the nature of the maximizing and minimizing vectors, for multiple roots.

4. The main result.

In Secs. 2 and 3 we have proved our result:

(4.1) Theorem. Given A , hermitian with eigenvalues $\{\lambda_i\}$, and b , arbitrary, define $\{b_i\}$ as in (1.3). Then the spectrum of (A, b) consists of all real λ such that

$$g(\lambda) = \sum_{i=1}^n \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda|^2} = 1 \quad (0/0 = 0; 1/0 = \infty) ,$$

together with each eigenvalue λ_k of A for which $g(\lambda_k) < 1$.

For each λ in the spectrum with $g(\lambda) = 1$, a unique x^λ is found by solving (1.7, 1.8). For each λ in the spectrum with $g(\lambda) < 1$, there exists an $(m-1)$ -sphere of x^λ satisfying (1.7, 1.8), where
 $m = \text{card } \{\lambda_j : \lambda_j = \lambda_k\}.$

Each x^λ so found renders $\Phi(x)$ stationary on S . Let
 $\Lambda_R = \max\{\lambda : \lambda \in \text{spectrum}\}$; let $\Lambda_L = \min\{\lambda : \lambda \in \text{spectrum}\}$. Let x_{\min}
 $= \text{any } x^{\Lambda_L}$; let $x_{\max} = \text{any } x^{\Lambda_R}$. Then $\Phi(x_{\min})$ minimizes $\Phi(x)$ on S ,
and $\Phi(x_{\max})$ maximizes $\Phi(x)$ on S .

5. The number of points in the spectrum.

As we noted in Sec. 2, if A is of order n , then the spectrum of (A, b) contains anywhere from 2 to $2n$ real numbers. When does it have the full amount $2n$? If any $\lambda_i b_i = 0$, then the discussion of Sec. 3 showed that the spectrum necessarily has fewer than $2n$ points. So we are limited to the case where all $\lambda_i b_i \neq 0$. But then, as shown in Sec. 2, we know that the spectrum is the intersection of the graph of

$$(5.1) \quad \mu = \sum_{i=1}^n \frac{\lambda_i^2 |b_i|^2}{|\lambda_i - \lambda|^2}$$

for real λ with the line $\mu = 1$.

The graph of (5.1) for real λ consists of $n + 1$ branches between the n vertical asymptotes $\lambda = \lambda_i$ ($i=1, \dots, n$). Since $\mu > 0$ for all λ , and $\mu \rightarrow 0$ as $\lambda \rightarrow \infty$ and $\lambda \rightarrow -\infty$, the right-most and left-most branches necessarily cut $\mu = 1$. The spectrum has the full number $2n$ of points if and only if each of the $n - 2$ interior branches of the curve reaches its minimum with $\mu < 1$. For general n a condition for this is probably too complicated to derive. For $n = 2$, however, we can answer the question, as follows:

(5.2) Theorem.

(5.3) Let $n = 2$, and assume A is in diagonal form with $\lambda_1 < \lambda_2$.
If the spectrum of (A, b) consists of 4 distinct numbers, then

$$(5.4) \quad 0 < |b_1 \lambda_1| \quad \text{and} \quad 0 < |b_2 \lambda_2| ,$$

and also

$$(5.5) \quad |b_1 \lambda_1|^{\frac{2}{3}} + |b_2 \lambda_2|^{\frac{2}{3}} < (\lambda_2 - \lambda_1)^{\frac{2}{3}} .$$

(5.6) Conversely, if (5.4) and (5.5) hold, then the spectrum of
 (A, b) consists of 4 distinct numbers.

Proof of (5.3). Let $a_i = |\lambda_i b_i|^2$ ($i=1,2$). If either a_1 or a_2 were zero, then the development in Sec. 3 shows that the spectrum would consist of at most 3 points. Hence $a_1 > 0$ and $a_2 > 0$; i.e., (5.4) holds. Let $M = (a_2/a_1)^{1/3}$. Now the development in Sec. 2 shows that the spectrum of (A, b) consists precisely of the real roots λ of the equation

$$(5.7) \quad g(\lambda) = \frac{a_1}{(\lambda - \lambda_1)^2} + \frac{a_2}{(\lambda - \lambda_2)^2} = 0 .$$

Since (5.7) has 4 real roots, we know that two roots must lie in the interval (λ_1, λ_2) . Now let μ be the unique real root of

$$g'(\lambda) = \frac{-2a_1}{(\lambda - \lambda_1)^3} - \frac{2a_2}{(\lambda - \lambda_2)^3} = 0 .$$

Then, because there are two roots of (5.7) in (λ_1, λ_2) ,

$$(5.8) \quad g(\mu) < 1 \quad .$$

We now show that (5.8) implies (5.4).

Solving $g'(\mu) = 0$ shows that

$$\frac{\lambda_2 - \mu}{\mu - \lambda_1} = M \quad ,$$

whence

$$\mu - \lambda_1 = \frac{1}{1+M} (\lambda_2 - \lambda_1) \quad ,$$

$$\lambda_2 - \mu = \frac{M}{1+M} (\lambda_2 - \lambda_1) \quad .$$

Hence

$$\begin{aligned} g(\mu) &= \frac{a_1(1+M)^2}{(\lambda_2 - \lambda_1)^2} + \frac{a_2(1+M)^2}{M^2(\lambda_2 - \lambda_1)^2} \\ &= \frac{(1+M)^2}{(\lambda_2 - \lambda_1)^2} \left[a_1 + \frac{a_2}{M^2} \right] \\ &= \frac{(1+M)^2 a_1^{\frac{2}{3}}}{(\lambda_2 - \lambda_1)^2} \left[\frac{1}{a_1^{\frac{1}{3}}} + \frac{1}{a_2^{\frac{1}{3}}} \right] \\ &= \frac{\left(\frac{1}{a_1^{\frac{1}{3}}} + \frac{1}{a_2^{\frac{1}{3}}} \right)^2}{(\lambda_2 - \lambda_1)^2} \left(\frac{1}{a_1^{\frac{1}{3}}} + \frac{1}{a_2^{\frac{1}{3}}} \right) \\ &= \frac{\left(\frac{1}{a_1^{\frac{1}{3}}} + \frac{1}{a_2^{\frac{1}{3}}} \right)^3}{(\lambda_2 - \lambda_1)^2} \quad . \end{aligned}$$

Thus $g(\mu) < 1$ implies

$$(5.9) \quad a_1^{\frac{1}{3}} + a_2^{\frac{1}{3}} < (\lambda_2 - \lambda_1)^{\frac{2}{3}},$$

which implies (5.5). Thus (5.3) is proved.

Proof of (5.6). We have $a_1 > 0$, $a_2 > 0$, and (5.9). The above steps are reversible, and so $g(\mu) < 1$, whence there are 4 real roots of $g(\mu) = 1$.

Thus theorem (5.2) is completely proved.

Condition (5.4) says that neither λ_1 nor λ_2 is 0, and that the point $b = (b_1, b_2)$ does not lie on an axis of the (x_1, x_2) -plane. Condition (5.5) requires that (b_1, b_2) be inside a curve Γ which depends only on the ratio λ_2/λ_1 . If $\lambda_2/\lambda_1 = 2$, for example, the curve Γ is $|b_1|^{2/3} + |2b_2|^{2/3} = 1$.




Figure 1

In Figure 1 the number of points in the spectrum of (A,b) is indicated for different b in the first quadrant by integers in circles.

If $|\lambda_2/\lambda_1| > 2$, the curve Γ includes values $|b_1| > 1$. But $|\lambda_2/\lambda_1| > 1$ implies that, on Γ , $|b_2| < 1$.

6. Geometrical interpretation.

The surfaces $\Phi(x) = k$ are similar conic surfaces with center b in the euclidean n -space \mathcal{E}_n of vectors x . The maximum problem (1.2) is to find the conic surface with maximum k which touches the constraint surface S , the unit sphere in \mathcal{E}_n . The rotation of A to diagonal form is a rotation of \mathcal{E}_n (leaving S invariant, of course) which causes principal axes of the conic surfaces to coincide with the axes of \mathcal{E}_n .

The vector $Ax - b$ is half the gradient of $\Phi(x)$, and x is the radius vector. Condition (1.5) merely states that at a point where $\Phi(x)$ is stationary, for x on S , the surface $\Phi(x) = k$ is tangent to S .

Fix x at a solution of (1.5), and let t be real. If the constant λ of (1.5) is positive, the value of $\Phi(tx)$ increases as t increases from 1; if λ is negative, $\Phi(tx)$ decreases as t increases from 1.

The main result of Secs. 2 and 3 is that the maximum problem of Sec. 1 is solved for the largest value of λ satisfying (1.5), for x on S . The authors see no obvious geometrical reason why this should be so.

If all $b_i \lambda_i \neq 0$, then Sec. 2 shows that any vector $x = x^\lambda$ which makes $\Phi(x)$ stationary on S is uniquely determined by λ .

Figure 2 shows, for $n = 2$ and $0 < \lambda_1 < \lambda_2$, a case where there are

4 distinct points of tangency of an ellipse with the unit circle. All ellipses have center b and common value of $\lambda_2/\lambda_1 > 2$. Since




Figure 2

$\lambda_2/\lambda_1 > 2$, it was shown in Sec. 5 that 4 distinct tangencies were possible for certain b outside S .

Whenever some $b_k = 0$, then, provided that (3.2) holds with the inequality sign $<$, we get more than one x belonging to a given λ . That is illustrated in Fig. 3, where $n = 2$ and $k = 1$. What is not obvious to the authors is a geometrical reason why necessarily $\lambda = \lambda_k$ in this case.




Figure 3

7. A constrained least squares problem.

Let C be an $m \times n$ matrix ($m \geq n$) and f an m -vector, both over the complex field. We wish to study the set of complex n -vectors y of euclidean length $\|y\| = (y^H y)^{1/2} = 1$ such that

$$(7.1) \quad \|Cy - f\|^2 = (Cy - f)^H (Cy - f) = \min.$$

The constraint is

$$(7.2) \quad \|y\|^2 = y^H y = 1.$$

Because euclidean length is invariant under unitary transformations, it is useful to rotate coordinates in both the space of y and the space of f . To do this, let $r = \text{rank}(C)$, and write

$$(7.3) \quad C = U^H D V,$$

where U, V are unitary, and where the only non-zero elements of D are the first r elements of the leading diagonal, which we may arrange so that

$$d_1 \geq d_2 \geq \cdots \geq d_r > 0.$$

Now let $V^H y = x$ and $Uf = g$. Then

$$\begin{aligned} Cy - f &= U^H D V y - U^H U f \\ &= U^H (Dx - g) \end{aligned}$$

Hence

$$\begin{aligned} \|Cy - f\|^2 &= \|Dx - g\|^2 \\ (7.4) \quad &= \sum_{i=1}^r |d_i x_i - g_i|^2 + \sum_{i=r+1}^m |g_i|^2 \end{aligned}$$

The problem (7.1, 7.2) is to minimize

$$\sum_{i=1}^r |d_i x_i - g_i|^2 = \sum_{i=1}^r d_i^2 |x_i - g_i/d_i|^2$$

subject to the constraint

$$(7.5) \quad \sum_{i=1}^n |x_i|^2 = 1$$

$$\text{Now let } \lambda_i = \begin{cases} 0 & (i = 1, 2, \dots, n-r) \\ d_{n+1-i}^2 & (i = n-r+1, \dots, n) \end{cases},$$

$$\text{and let } b_i = \begin{cases} 0 & (i = 1, 2, \dots, n-r) \\ \frac{g_{n+1-i}}{d_{n+1-i}} & (i = n-r+1, \dots, n) \end{cases}.$$

We then have changed our problem to one of minimizing

$$(7.6) \quad \sum_{i=1}^n \lambda_i |x_i - b_i|^2, \quad ,$$

subject to the constraint (7.5), where

$$(7.7) \quad 0 = \lambda_1 = \dots = \lambda_{n-r} < \lambda_{n-r+1} \leq \dots \leq \lambda_n .$$

This is precisely the minimum problem (1.2) of Sec. 1. The special role of the $n - r$ zero eigenvalues of $C^H C$ becomes evident.

Thus the general problem of the least-squares solution of $Cy = f$ with constraint (7.2) is a special case of our minimum problem (1.2).

8. Lemmas from complex function theory.

In this final section we state and prove three lemmas relating partial derivatives of certain regular analytic functions of several complex variables to gradients of real-valued functions of vector variables. This technique is common in the study of second-order partial differential equations; for example, see (9.1). We include the material mainly to keep our treatment self-contained, and partly to call explicit attention to the fact that the Lagrange multiplier λ must be real even though complex variables are used.

(8.1) Lemma. Let $\Phi(\lambda, \mu)$ be a regular analytic function of two complex variables λ, μ such that, for all real x, y ,

$$(8.2) \quad F(x,y) = \Phi(x+iy, x-iy)$$

is real-valued. Then

$$\frac{\partial F}{\partial x} + i \frac{\partial F}{\partial y} = 2 \left. \frac{\partial \Phi}{\partial \mu} \right|_{\substack{\lambda=x+iy \\ \mu=x-iy}} .$$

Proof. Differentiate (8.2):

$$(8.3) \quad \frac{\partial F}{\partial x} = \frac{\partial \Phi}{\partial \lambda} \cdot 1 + \frac{\partial \Phi}{\partial \mu} \cdot 1 ,$$

$$(8.4) \quad \frac{\partial F}{\partial y} = \frac{\partial \Phi}{\partial \lambda} \cdot i - \frac{\partial \Phi}{\partial \mu} \cdot i .$$

Add (8.3) to (8.4) $\times i$:

$$\frac{\partial F}{\partial x} + i \frac{\partial F}{\partial y} = 2 \frac{\partial \Phi}{\partial \mu} .$$

(8.5) Lemma. Let F and G be real-valued differentiable functions of real variables $x_1, y_1, \dots, x_n, y_n$. For abbreviation, let $z_k = x_k + iy_k$, and let $z = (z_1, \dots, z_n)$. Then, for $F(z)$ to be stationary at $z = a$ with respect to all neighboring z such that $G(z) = G(a)$, it is necessary and sufficient that there exist a real Lagrange constant λ such that

$$(8.6) \quad \frac{\partial F}{\partial x_k} + i \frac{\partial F}{\partial y_k} - \lambda \left(\frac{\partial G}{\partial x_k} + i \frac{\partial G}{\partial y_k} \right) = 0$$

for $z = a$ and $k = 1, \dots, n$.

Proof. Condition (8.6) is nothing but the usual condition that the real gradient vector

$$\left(\frac{\partial F}{\partial x_1}, \frac{\partial F}{\partial y_1}, \frac{\partial F}{\partial x_2}, \frac{\partial F}{\partial y_2}, \dots, \frac{\partial F}{\partial x_n}, \frac{\partial F}{\partial y_n} \right)$$

be parallel to the vector

$$\left(\frac{\partial G}{\partial x_1}, \frac{\partial G}{\partial y_1}, \frac{\partial G}{\partial x_2}, \frac{\partial G}{\partial y_2}, \dots, \frac{\partial G}{\partial x_n}, \frac{\partial G}{\partial y_n} \right).$$

The use of the complex variables z_k is unessential.

Given any vector $z = (z_1, \dots, z_n)$, we let \bar{z} denote the vector of complex conjugates $(\bar{z}_1, \dots, \bar{z}_n)$.

(8.7) Lemma. Let $\Phi(z, w)$ and $\Psi(z, w)$ be regular analytic functions of the two complex vector variables $z = (z_1, \dots, z_n)$ and $w = (w_1, \dots, w_n)$ with the property that $\Phi(z, \bar{z})$ and $\Psi(z, \bar{z})$ are real. Then $\Phi(z, \bar{z})$ is stationary at $z = a$ with respect to all z such that $\Psi(z, \bar{z}) = \Psi(a, \bar{a})$, if and only if there exists a real Lagrange constant λ such that

$$\frac{\partial \Phi}{\partial w_k} - \lambda \frac{\partial \Psi}{\partial w_k} = 0$$

for $z = a$ and $w = \bar{a}$ and $k = 1, 2, \dots, n$.

Proof. Let $z = x + iy$. Then $\Phi(z, \bar{z}) = F(x, y)$, $\Psi(z, \bar{z}) = G(x, y)$. By lemma 8.1 applied to each variable z_k ,

$$\frac{\partial \Phi}{\partial w_k} = \frac{\partial F}{\partial x_k} + i \frac{\partial F}{\partial y_k} ,$$

$$\frac{\partial \Psi}{\partial w_k} = \frac{\partial G}{\partial x_k} + i \frac{\partial G}{\partial y_k}$$

for $z = a$, $w = \bar{a}$, and $k = 1, \dots, n$.

Then lemma (8.7) follows from lemma (8.5) .

9. References.

- (9.1) P. R. Garabedian, Partial Differential Equations, Wiley, 1964; Chap. 3.
- (9.2) David L. Phillips, A technique for the numerical solution of certain integral equations of the first kind, J. Assoc. Comput. Mach. vol. 9 (1962), pp. 84-97.
- (9.3) S. Twomey, On the numerical solution of Fredholm integral equations of the first kind by the inversion of the linear system produced by quadrature, J. Assoc. Comput. Mach. vol. 10 (1963), pp. 97-101.

STANFORD UNIVERSITY
TECHNICAL REPORTS DISTRIBUTION LIST
CONTRACT Nore-225(37)
(NR 044-211)

ARMED SERVICES TECHNICAL INFORMATION AGENCY ARLINGTON HALL STATION ARLINGTON 12, VIRGINIA	10	OFFICE OF TECHNICAL SERVICES DEPARTMENT OF COMMERCE WASHINGTON 25, D. C.	1	PROF. GEORGE W. BROWN WESTERN DATA PROCESSING CENTER UNIVERSITY OF CALIF. LOS ANGELES 24, CALIF.	1
CHIEF OF NAVAL RESEARCH CODE 432 OFFICE OF NAVAL RESEARCH WASHINGTON 25, D.C.	2	DOCUMENT CENTER STANFORD RESEARCH INSTITUTE 333 RAVENSWOOD AVE. MENLO PARK, CALIF.	1	PROF. JOHN W. CARR, III MOORE SCHOOL OF E. E. ENGR. UNIVERSITY OF PENNSYLVANIA PHILADELPHIA 4, PENN.	1
COMMANDING OFFICER OFFICE OF NAVAL RESEARCH BRANCH OFFICE 1000 GEARY ST. SAN FRANCISCO 9, CALIF.	1	TECHNICAL INFORMATION OFFICER NAVAL RESEARCH LABORATORY WASHINGTON 25, D. C.	6	MR. ROBERT L. CAUSEY LOCKHEED AIRCRAFT CORP. MISSILES AND SPACE 3251 HANOVER ST. PALO ALTO, CALIF.	1
COMMANDING OFFICER OFFICE OF NAVAL RESEARCH BRANCH OFFICE 495 SUMMER STREET BOSTON 10, MASSACHUSETTS	1	THE MITRE CORP. LIBRARY ATTN. SUPERVISOR LIBRARY SERVICES BEDFORD, MASSACHUSETTS	2	PROF. ABRAHAM CHARNES THE TECHNOLOGICAL INSTITUTE NORTHWESTERN UNIVERSITY EVANSTON, ILLINOIS	1
COMMANDING OFFICER OFFICE OF NAVAL RESEARCH BRANCH OFFICE 346 BROADWAY NEW YORK 13, NEW YORK	1	UNIVERSITY OF CALIFORNIA ATTN. LIBRARY, NUMERICAL ANALYSIS RES. LOS ANGELES 24, CALIF.	1	PROF. B. A. CHARTRES DIVISION OF APPLIED MATHEMATICS BROWN UNIVERSITY PROVIDENCE 12, RHODE ISLAND	1
COMMANDING OFFICER OFFICE OF NAVAL RESEARCH BRANCH OFFICE NAVY NO. 100, FLEET POST OFFICE NEW YORK, NEW YORK	2	UNIVERSITY OF MIAMI SCIENTIFIC COMPUTING CENTER P.O. BOX 8011 CORAL GABLES 46, FLORIDA	1	PROF. HARVEY COHN DEPARTMENT OF MATHEMATICS UNIVERSITY OF ARIZONA TUSCON 25, ARIZONA	1
COMMANDING OFFICER OFFICE OF NAVAL RESEARCH BRANCH OFFICE 1030 EAST GREEN ST. PASADENA 1, CALIF.	1	U.S. NAVAL ORDNANCE LABORATORY WHITE OAK ATTN: MATHEMATICS DEPARTMENT SILVER SPRING, MARYLAND	1	PROF. STEPHEN CRANDALL DEPARTMENT OF MECHANICS MASS. INST. OF TECHNOLOGY CAMBRIDGE 39, MASSACHUSETTS	1
ARGONNE NATIONAL LABORATORY TECHNICAL INFORMATION DIVISION 9700 SOUTH CASS AVE. ARGONNE, ILLINOIS	1	U. S. NAVAL ORDNANCE LABORATORY WHITE OAK ATTN. LIBRARY SILVER SPRING, MARYLAND	1	PROF. PHILIP DAVIS DEPARTMENT OF MATHEMATICS BROWN UNIVERSITY PROVIDENCE 12, RHODE ISLAND	1
COMMANDING OFFICER U.S. NAVAL ELECTRONICS LAB. ATTN. R. F. ARENZ F. A. SABRANSKY SAN DIEGO 52, CALIF.	2	U. S. NAVY MANAGEMENT OFFICE DEPARTMENT OF THE NAVY WASHINGTON 25, D. C.	1	PROF. DAVID B. DEKKER DIRECTOR, RESEARCH COMPUTER LAB. UNIVERSITY OF WASHINGTON SEATTLE 5, WASHINGTON	1
COMPUTATION CENTER UNIVERSITY OF TEXAS AUSTIN 12, TEXAS	1	U. S. NAVAL WEAPONS LABORATORY DAHLGREN, VIRGINIA	1	PROF. R. J. DEVOGELAERE DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA BERKELEY 4, CALIF.	1
COMPUTER REVIEWS ROOM 5-C 211 E. 43rd STREET NEW YORK 36, NEW YORK	1	PROFESSOR W. F. ATCHISON RICH ELECTRONIC COMPUTER CENTER GEORGIA INSTITUTE OF TECHNOLOGY ATLANTA, GEORGIA	1	PROF. C. L. DOLPH 311 PARK AVENUE ANN ARBOR, MICHIGAN	1
DEPARTMENT OF MATHEMATICS HARVARD UNIVERSITY CAMBRIDGE, MASSACHUSETTS	1	DR. ERWIN H. BAREISS APPLIED MATHEMATICS DIVISION ARGONNE NATIONAL LABORATORY 9700 SOUTH CASS AVE. ARGONNE, ILLINOIS	1	PROF. JIM DOUGLAS, JR. DEPARTMENT OF MATHEMATICS THE RICE INSTITUTE HOUSTON 1, TEXAS	1
DEPARTMENT OF MATHEMATICS MASSACHUSETTS INST. OF TECH CAMBRIDGE, MASSACHUSETTS	1	PROF. ROBERT C. F. BARTELS COMPUTING CENTER UNIVERSITY OF MICHIGAN ANN ARBOR, MICHIGAN	1	PROF. R. J. DUFFIN MATHEMATICS DEPARTMENT CARNEGIE INSTITUTE OF TECHNOLOGY PITTSBURGH 13, PENNSYLVANIA	1
SPACE TECHNOLOGY LABS. ONE SPACE PARK ATTN. IRENE MATHEWS REDONDO BEACH, CALIF.	1	DR. RICHARD BELLMAN THE RAND CORPORATION 1700 MAIN ST. SANTA MONICA, CALIFORNIA	1	DR. ROBERT DICKSON LOCKHEED AIRCRAFT CORPORATION MISSILES AND SPACE DIVISION 3251 HANOVER STREET PALO ALTO, CALIF.	1
INSTITUTE FOR MATHEMATICAL SCIENCES NEW YORK UNIVERSITY NEW YORK, 3, NEW YORK	1	PROF. GARRETT BIRKHOFF MATHEMATICS DEPARTMENT HARVARD UNIVERSITY CAMBRIDGE 38, MASSACHUSETTS	1	PROF. PAUL DWYER 403 LENAWEE DRIVE ANN ARBOR, MICHIGAN	1
LIBRARY DIGITAL COMPUTER LAB. UNIVERSITY OF ILLINOIS URBANA, ILLINOIS	1	BERNARD H. BISSINGER DEPARTMENT OF MATHEMATICS MATH LIBRARY LEBANON VALLEY COLLEGE ANNVILLE, PENNSYLVANIA	1	MR. THOMAS A. DWYER DIRECTOR, COMPUTER SCIENCE DEPARTMENT OF PHYSICS UNIVERSITY OF DAYTON DAYTON 9, OHIO	1
LIBRARIAN, COMMUNICATIONS RESEARCH DIVISION INSTITUTE FOR DEFENCE ANALYSES VON NEUMANN HALL PRINCETON, NEW JERSEY	1	MR. TRUMAN BOTTS DEPARTMENT OF MATHEMATICS UNIVERSITY OF VIRGINIA CHARLOTTESVILLE, VIRGINIA	1	MR. EDGAR L. EICHNORH BURROUGHS CORPORATION ELECTRADATA FACILITY 460 SIERRA MADRE VILLA PASADENA, CALIF.	1
NATIONAL AERONAUTICS AND SPACE ADM. LEWIS RESEARCH CENTER 21000 BROOKPARK ROAD ATTN. LIBRARY CLEVELAND 35, OHIO	1	PROF. JAMES BRAMBLE INSTITUTE FLUID DYNAMICS AND APPLIED MATHEMATICS UNIVERSITY OF MARYLAND COLLEGE PARK, MARYLAND	1	MR. FRANK ENGEL, JR. ANALYTICAL DEPARTMENT 4-L-39 WESTINGHOUSE ELECTRIC CORP. 700 BRADDOCK AVENUE EAST PITTSBURGH, PENNSYLVANIA	1

DR. G. W. EVANS, II STANFORD RESEARCH INSTITUTE 333 RAVENSWOOD AVENUE MENLO PARK, CALIF.	1	PROF. B. E. HUBBARD INSTITUTE FLUID DYNAMICS AND APPLIED MATHEMATICS UNIVERSITY OF MARYLAND COLLEGE PARK, MARYLAND	1	B. M. PARLETT INST. FOR MATHEMATICAL SCIENCES 4 WASHINGTON PLACE NEW YORK 3, NEW YORK	1
PROF. F. A. FICKEN DEPARTMENT OF MATHEMATICS NEW YORK UNIVERSITY UNIVERSITY HEIGHTS NEW YORK 53, NEW YORK	1	DR. M. A. HYMAN IBM SYSTEMS DEVELOPMENT CENTER 7220 WISCONSIN AVE. BETHESDA, MARYLAND	1	DR. SEYMOUR V. PARTER DEPARTMENT OF MATHEMATICS UNIVERSITY OF WISCONSIN MADISON, WISCONSIN	1
MR. WERNER FRANK 4617 LOARKWOOD AVENUE WOODLAND HILLS, CALIF.	1	PROF. EUGENE ISAACSON INSTITUTE OF MATHEMATICAL SCIEN. NEW YORK UNIVERSITY 4 WASHINGTON PLACE NEW YORK 3, NEW YORK	1	PROF. ALAN J. PERLIS COMPUTATION CENTER CARNEGIE INSTITUTE OF TECHNOLOGY PITTSBURGH 13, PENNSYLVANIA	1
PROF. J. N. FRANKLIN CALIFORNIA INST. OF TECHNOLOGY PASADENA 4, CALIF.	1	DR. W. J. JAMESON RESEARCH DIVISION COLLINS RADIO COMPANY CEDAR RAPIDS, IOWA	1	PROF. CLAY L. PERRY DIRECTOR, COMPUTATION CENTER UNIVERSITY OF CALIFORNIA SCHOOL OF SCIENCE AND ENGINEERING LA JOLLA, CALIF.	1
PROF. BERNARD FRIEDMAN DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA BERKELEY 4, CALIF.	1	PROF. M. KATZ DEPARTMENT OF MATHEMATICS UNIVERSITY OF WISCONSIN MILWAUKEE 11, WISCONSIN	1	PROF. MURRAY I. TER DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA BERKELEY 4, CALIF.	1
DR. H. H. GERMOND RCA SERVICE COMPANY BLDG. 989-1 MU 811 PATRICK AIR FORCE BASE FLORIDA	1	PROF. THOMAS A. KEEMAN COMPUTING CENTER UNIVERSITY OF ROCHESTER ROCHESTER 20, NEW YORK	1	PROF. WERNER RHEINBOLDT COMPUTER SCIENCE CENTER UNIVERSITY OF MARYLAND COLLEGE PARK, MARYLAND	1
PROF. WALLACE GIVENS DEPARTMENT OF ENGR. SCIENCE NORTHWESTERN UNIVERSITY EVANSTON, ILLINOIS	1	PROF. WILLIAM B. KEHL COMPUTATION AND DATA PROC. CENT. 825 CATHEDRAL OF LEARNING UNIVERSITY OF PITTSBURGH PITTSBURGH 13, PENNSYLVANIA	1	PROF. R. D. RICHTMYER AEC COMPUTING AND APPLIED MATHEMATICS CENTER INSTITUTE OF MATHEMATICAL SCIENCES NEW YORK UNIVERSITY 4 WASHINGTON PLACE NEW YORK 3, NEW YORK	1
DR. H. M. GOLDSTINE RESEARCH CENTER INTERNATIONAL BUSINESS MACHINES YORKTOWN, NEW YORK	1	DR. ERYAND G. KOGDETLIANTZ ROCKEFELLER INST. FOR MEDICAL RESEARCH 66th ST. AND YORK AVE. NEW YORK, NEW YORK	1	MRS. MAXINE ROCKHOFF DEPARTMENT OF COMMERCE NATIONAL BUREAU OF STANDARDS WASHINGTON 25, D. C.	1
PROF. SAUL GORN MOORE SCHOOL OF ELEC. ENGINEERING UNIVERSITY OF PENNSYLVANIA PHILADELPHIA 4, PENNSYLVANIA	1	PROF. R. E. LANGER DEPARTMENT OF MATHEMATICS UNIVERSITY OF WISCONSIN MADISON 6, WISCONSIN	1	EDWARD M. ROSEN SENIOR MATHEMATICIAN APPLIED MATHEMATICS DEPARTMENT MONSANTO CHEMICAL CO. 800 N. LINDBERGH ST. LOUIS 66, MISSOURI	1
PROF. L. GREENBERG DEPARTMENT OF MATHEMATICS BROWN UNIVERSITY PROVIDENCE, RHODE ISLAND	1	PROF. RALPH E. LEE DIRECTOR OF COMPUTER CENTER SCHOOL OF MINES AND METALLURGY UNIVERSITY OF MISSOURI ROLLA, MISSOURI	1	DR. SAMUEL SCHECHTER RESEARCH SCIENTIST AEC COMPUTING CENTER NEW YORK UNIVERSITY NEW YORK 3, NEW YORK	1
MR. SAMUEL W. GREENHOUSE CHIEF, SECTION OF STAT. AND MATH NATIONAL INSTITUTE OF MENTAL HEALTH BETHESDA 14, MARYLAND	1	PROF. ARVID LONSETH DEPARTMENT OF MATHEMATICS OREGON STATE COLLEGE CORVALLIS, OREGON	1	PROF. LOWELL SCHOENFELD MATHEMATICS DEPARTMENT PENNSYLVANIA STATE UNIVERSITY STATE COLLEGE, PENNSYLVANIA	1
PROF. R. T. GREGORY COMPUTATION CENTER UNIVERSITY OF TEXAS AUSTIN 12, TEXAS	1	MR. B. MITTMAN ARMOUR RESEARCH FOUNDATION 10 WEST 35TH STREET CHICAGO 16, ILLINOIS	1	DR. DANIEL SHANKS DAVID TAYLOR MODEL BASIN WASHINGTON 7, D. C.	1
DR. T. M. E. GREVILLE MATH. RESEARCH CENTER, U.S. ARMY UNIVERSITY OF WISCONSIN MADISON 6, WISCONSIN 53706	1	PROF. JAMES M. MOORE DEPARTMENT OF INDUSTRIAL ENGINEERING NORTHEASTERN UNIVERSITY BOSTON 12, MASSACHUSETTS	1	JAMES M. SLOSS MATHEMATICS DEPARTMENT UNIVERSITY OF CALIFORNIA SANTA BARBARA, CALIF.	1
PROF. PRESTON HAMMER DIRECTOR, NUMERICAL ANALYSIS LAB. UNIVERSITY OF WISCONSIN MADISON 6, WISCONSIN	1	PROF. T. S. MOTZKIN DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA BERKELEY 4, CALIF.	1	PROF. THOMAS H. SOUTHARD CHAIRMAN, MATHEMATICS DEPARTMENT ALAMEDA STATE COLLEGE 22300 FOOTHILL BLVD. HAYWARD, CALIFORNIA	1
DR. ELDON HANSEN 347 ALDEAN MOUNTAIN VIEW, CALIF.	1	DR. J. P. NASH LOCKHEED MISSILE AND SPACE CORP. SUNNYVALE, CALIF.	1	PROF. M. L. STEIN DEPARTMENT OF MATHEMATICS INSTITUTE OF TECHNOLOGY UNIVERSITY OF MINNESOTA MINNEAPOLIS 4, MINNESOTA	1
PROF. T. J. HIGGINS DIRECTOR, UNIVERSITY OF WISCONSIN ENGINEERING EXPERIMENT STATION MADISON 6, WISCONSIN	1	PROF. C. D. OLDS P. O. BOX 462 LOS ALTOS, CALIFORNIA	1	PROF. J. J. STOKER INSTITUTE OF MATHEMATICAL SCIENCES 25 WAVERLY PLACE NEW YORK 3, NEW YORK	1
PROF. T. W. HILDEBRANDT DEPARTMENT OF MATHEMATICS OHIO STATE UNIVERSITY COLUMBUS, OHIO	1	DR. JAMES M. ORTEGA DIVISION 5426 SANDIA CORPORATION ALBUQUERQUE, NEW MEXICO	1	MR. H. C. THACHER, JR. ARGONNE NATIONAL LABORATORY 9700 SOUTH CASS AVENUE ARGONNE, ILLINOIS	1
DR. WILLIAM HOOKER PHILCO CORPORATION 3875 FABIAN WAY PALO ALTO, CALIF.	1	DR. JOSEPH OTTERMAN P. O. BOX 2008 UNIVERSITY OF MICHIGAN ANN ARBOR, MICHIGAN	1		
DR. A. S. HOUSEHOLDER OAK RIDGE NATIONAL LAB. P. O. BOX X OAK RIDGE, TENNESSEE	1				

MR. ROLAND THOMPSON PURROUGHS CORPORATION 6071 2ND AVE. DETROIT 32, MICHIGAN	1	PROF. DR. L. COLLATZ INSTITUT FUR ANGEWANDTE MATH UNIVERSITAT HAMBURG (24A) HAMBURG 13 ROTHENBAUMCHAUSSÉE 67/69 GERMANY	1	PROF. E. S. PAGE DIRECTOR, UNIVERSITY COMPUTING LAB. NEWCASTLE ON TYNE ENGLAND	1
DR. GENE THOMPSON UNITED TECHNOLOGY CORP. 1050 E. ARQUES AVE. SUNNYVALE, CALIF.	1	PROF. GERMUND G. DAHLQVIST ROYAL INST. OF TECHNOLOGY STOCKHOLM DIV. OF APPLIED MATH STOCKHOLM 70, SWEDEN	1	PROF. DR. R. SAUER MATHEMATISCHES INSTITUT DER TECHNISCHEN HOCHSCHULE MÜNCHEN 2 ARCSSTR. 21, GERMANY	1
PROF. JOHN TODD MATHEMATICS DEPARTMENT CALIF. INST. OF TECHNOLOGY PASADENA 4, CALIFORNIA	1	DR. LESLIE FOX OXFORD UNIVERSITY OXFORD, ENGLAND	1	PROF. E. STIEFEL INSTITUT FUR ANGEWANDTE MATH. EIDG. TECHNISCHE HOCHSCHULE ZÜRICH, SWITZERLAND	1
PROF. RICHARD S. VARGA DEPARTMENT OF MATHEMATICS CASE INSTITUTE OF TECHNOLOGY UNIVERSITY CIRCLE CLEVELAND 6, OHIO	1	DR. CARL-ERIK FROBERG ADV. FOR NUMERISK ANALYSIS LUNDS UNIVERSITET LUND, SWEDEN	1	PROF. DR. HEINZ UNGER INSTITUT FUR ANGEWANDTE MATH. DER UNIVERSITAT BONN WEGELERSTRASSE 10 BONN, GERMANY	1
PROF. WOLFGANG WASOW DEPARTMENT OF MATHEMATICS UNIVERSITY OF WISCONSIN MADISON 6, WISCONSIN	1	MR. RICHARD GOODMAN DIRECTOR, AUTOMATIC PROGRAMMING INFORMATION CENTER DEPARTMENT OF MATHEMATICS THE TECHNICAL COLLEGE BRIGHTON, ENGLAND	1	PROF. DR. A. WALTHER DIREKTOR, INSTITUT FUR PRAKTISCHE MATHEMATIK TECHNISCHE HOCHSCHULE (16) DARMSTADT, GERMANY	1
PROF. M. G. WURTELE DEPARTMENT OF METEOROLOGY UNIVERSITY OF CALIFORNIA LOS ANGELES 24, CALIF.	1	DR. J. M. HAMMERSLEY INSTITUTE OF STATISTICS ST. CROSS ROAD OXFORD, ENGLAND	1	DR. S. B. L. WILSON COMPUTING DEPARTMENT AERO ENGINE DIVISION ROLLS-ROYCE LIMITED ELTON ROAD DERBY, ENGLAND	1
PROF. DAVID M. YOUNG, JR. DEPARTMENT OF MATHEMATICS UNIVERSITY OF TEXAS AUSTIN 12, TEXAS	1	PROF. DR. PETER HENRICI LEHRSTUHL FUR HOHERE MATHEMATIK EIGENOSSENSISCHE TECHNISCHE HOCHSCHULE ZÜRICH, SWITZERLAND	1	PROF. RIGAL SECRETAIRE A L'AFCAI 48 RUE DES POTIERS MONTBELIARD, DOUBS FRANCE	1
<u>DISTRIBUTION VIA ONR LONDON</u>					
F. C. ADEY F. L. A., CHIEF LIBRARIAN LEICESTER COLLEGE OF ART AND TECH. THE NEWARKS, LEICESTER, ENGLAND	1	DR. JOSEPH HERSCH E. T. H. ZIMMER 200 ZÜRICH 6, SWITZERLAND	1	PROF. DR. H. W. WIELANDT MATHEMATISCHES INSTITUT UNIVERSITAT TUBINGEN TUBINGEN (14B) WILHELMSTRASSE 7, GERMANY	1
MATHEMATISCH CENTRUM 2E BOLRIJAAVESTRAT 49 AMSTERDAM 0, NETHERLANDS	1	INTERNATIONAL COMPUTATION CENTER PALAZZO DEGLI UFFICI ZONA DELL'E. U. R. ROMA, ITALY	1	DR. M. V. WILKES UNIVERSITY MATHEMATICAL LAB. CORN EXCHANGE STREET CAMBRIDGE, ENGLAND	1
PROF. A. C. AITKEN MATHEMATICAL INSTITUTE UNIVERSITY OF EDINBURGH 16 CHAMBERS STREET EDINBURGH 1, SCOTLAND	1	PROF. J. KUNTZMANN LABORATOIRE DE CALCUL UNIVERSITE DE GRENOBLE 44-46 AVENUE FELIX-VIALLET GRENOBLE, ISERE FRANCE	1	DR. J. H. WILKINSON MATHEMATICS DIVISION NATIONAL PHYSICAL LAB. TEDDINGTON, MIDDLESEX ENGLAND	1
DR. ENZO APARO ISTITUTO NAZIONALE PER LE APPLICAZIONI DEL CALCOLO PIAZZALE DUE SCIENZE 7 ROMA, ITALY	1	PROF. PENTTI LAASONEN FINLAND INSTITUTE OF TECHNOLOGY HELSINKI, FINLAND	1	HONORARY LIBRARIAN, BRITISH COMPUTER SOCIETY, FINSBURY COURT, FINSBURY PAVEMENT LONDON E. C. 2, ENGLAND	1
PROF. DR. CHARLES BLANC CENTRE DE CALCUL ELECTRONIQUE AVENUE DE COUR, 33 LAUSANNE, SWITZERLAND	1	PROF. C. LANCZOS INSTITUTE FOR ADVANCED STUDIES 64-65 MERRION SQUARE DUBLIN, IRELAND	1	THE LIBRARY, BUILDING 465 ATOMIC ENERGY RES. ESTABLISHMENT HARWELL, DIDCOT BERKSHIRE, ENGLAND	1
JEAN A. BAUDOT COMPUTING CENTRE DEPARTMENT OF MATHEMATICS UNIVERSITE DE MONTREAL MONTREAL, CANADA	1	PROF. J. L. LIONS DEPARTMENT DE MATHEMATIQUES UNIVERSITE DE PARIS 11 RUE PIERRE CURIE PARIS 5E, FRANCE	1	JEAN A. BAUDOT COMPUTING CENTER DEPARTMENT OF MATHEMATICS UNIVERSITE DE MONTREAL MONTREAL, CANADA	1
PROF. DR. F. L. BAUER MATHEMATISCHES INST. DER TECHNISCHEN HOCHSCHULE 8 MÜNCHEN 2 ARCSSTRASSE 21 GERMANY	1	PROF. L. MIRSKY THE UNIVERSITY SHEFFIELD, ENGLAND	1	<u>OTHER FOREIGN ADDRESSES</u>	
MR. NIELS IVAR BECH REGNECENTRALEN G1. CARLSBERGVEJ 2 VALBY, DENMARK	1	DR. A. R. MITCHELL MATHEMATICS DEPARTMENT ST. SALVATORS COLLEGE UNIVERSITY OF ST. ANDREWS ST. ANDREWS, FIFE, SCOTLAND	1	DR. JOHN M. BENNETT ADOLPH BASSER COMPUTER LAB. UNIVERSITY OF SYDNEY SYDNEY, N.S.W. AUSTRALIA	1
J. G. BYRNE ENGINEERING SCHOOL TRINITY COLLEGE DUBLIN 2, IRELAND	1	MR. JEAN A. MUSSARD EXECUTIVE SECRETARY PROVISIONAL INTERNATIONAL COMPUTATION CENTER PALAZZO DEGLI UFFICI ZONA DELL'E. U. R. ROMA, ITALY	1	W. A. COWAN, LIBRARIAN BARR SMITH LIBRARY THE UNIVERSITY ADELAIDE SOUTH AUSTRALIA	1
DR. STIG COMET INTERNATIONAL COMPUTATION CENTRE PALAZZO DEGLI UFFICI ZONA DELL'E. U. R. ROMA, ITALY	1	PROF. A. OSTROWSKI CERTENAGO-MONTAGNOLA BEL LUGANO SWITZERLAND		PROF. A. GONZALEZ DOMINGUEZ THE INSTITUTE OF CALCULUS UNIVERSITY OF BUENOS AIRES A' ENIDA DE MAYO 760, 20 PISO BUENOS AIRES, ARGENTINA	1

FACULTAD DE CIENCIAS EXACTAS
Y NATURAL
BIBLIOTECA SECCION CANJE
CASILLA DE CORREO 1766
BUENOS AIRES
REPUBLICA ARGENTINA

1

PROF. SANTIAGO FRIEDMANN
DIRECTOR DEL CENTRO DE COMPUTACION
UNIVERSIDAD DE CHILE
CASILLA 2777
SANTIAGO, CHILE

1

PROF. ALAN GIBB
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ALBERTA
CALBARY, ALBERTA
CANADA

1

DR. C. C. GOTLIEB
COMPUTATION CENTER
MCLENNAN LABORATORY
UNIVERSITY OF TORONTO
TORONTO, CANADA

1

DR. J. F. HART
HEAD, COMPUTING CENTRE
THE UNIVERSITY OF WESTERN ONTARIO
LONDON, CANADA

1

PROF. THOMAS E. HULL
COMPUTATION CENTER
MCLENNAN LABORATORY
UNIVERSITY OF TORONTO
TORONTO, CANADA

1

PROF. W. KAHAN
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TORONTO
TORONTO, CANADA

1

PROF. J. MCNAMEE
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ALBERTA
ALBERTA, CANADA

1

PROF. MANUEL SADOSKY
INSTITUTO DE CALCULO
FACULTAD DE CIENCIAS EXACTAS
Y NATURALES
PERU 27, BUENOS AIRES
REPUBLICA ARGENTINA

1

PROF. H. SCHWERTFEGER
DEPARTMENT OF MATHEMATICS
MCGILL UNIVERSITY
MONTREAL, QUEBEC
CANADA

1

MR. J. C. VELLA
PUBLISHING SECRETARY
SOCIEDAD ARGENTINA DE CALCULO
AVENIDO DE MAYO 760 - 20 PISO
BUENOS AIRES,
ARGENTINA

2

ADDITIONAL COPIES FOR FUTURE
REQUIREMENTS

50

CONTRACT NONR-225(37)
NOVEMBER 1963
247

PROF. R. KODIS
DEPARTMENT OF MATHEMATICS
BROWN UNIVERSITY
PROVIDENCE, RHODE ISLAND

1

PROF. C. DONALD LA BUDE
DEPARTMENT OF CHEMISTRY
NEW YORK UNIVERSITY
UNIVERSITY HEIGHTS
NEW YORK 53, NEW YORK

1

UNCLASS

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Computer Science Department Stanford University Stanford, Calif. 94305		2a. REPORT SECURITY CLASSIFICATION UNCLASS
		2b. GROUP -----
3. REPORT TITLE MAXIMIZING A SECOND-DEGREE POLYNOMIAL ON THE UNIT SPHERE		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Manuscript for Publication (Technical Report)		
5. AUTHOR(S) (Last name, first name, initial) FORSYTHE, George E. and GOLUB, Gene H.		
6. REPORT DATE February 5, 1965	7a. TOTAL NO. OF PAGES 31	7b. NO. OF REFS 3
8a. CONTRACT OR GRANT NO. Nonr-225(37)	8b. ORIGINATOR'S REPORT NUMBER(S) CS16	
9. PROJECT NO. NR-044-211		
	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) none	
10. AVAILABILITY/LIMITATION NOTICES 1. Qualified requesters may obtain copies of this report from: Defence Document Center, Cameron Station Alexandria, Virginia. 2. Chief, Input Section, Clearinghouse for Federal Scientific & Tech. Information, CFSTI Sills Bldg. 5285 Port Royal Road, Springfield, Virginia 22151.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY OFFICE OF NAVAL RESEARCH Code 432	
13. ABSTRACT Let A be a hermitian matrix of order n , and b a known vector in C^n . The problem is to determine which vectors make $\phi(x) = (x-b)^H A(x-b)$ a maximum or minimum on the unit sphere $U = \{x : x^H x = 1\}$. The problem is reduced to the determination of a finite point set, the spectrum of (A,b) . The theory reduces to the usual theory of hermitian forms when $b = 0$.		

DD FORM 1473
JAN 64

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
1. Maximization on a sphere						
2. Minimization on a sphere						
3. Constrained maximization-minimization						
4. Maximization of quadratic function						
5. Minimization of quadratic function						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive S200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

DD FORM 1473 (BACK)

1 JAN 64

UNCLASSIFIED

Security Classification